

英文原著・総説・症例報告

著者	タイトル	掲載誌名等
Abulimiti A, Nishitani-Yokoyama M, Shimada K, Kunimoto M, Matsubara T, et al.	Prognostic impact of peak oxygen uptake and heart rate reserve in patients after off-pump coronary artery bypass grafting.	Clin Cardiol. 2021;44(4):580-587.
Aizawa J, Hirohata K, Ohji S, Ohmi T, Mitomo S, et al.	Correlations between isokinetic knee torques and single-leg hop distances in three directions in patients after ACL reconstruction.	BMC Sports Sci Med Rehabil. 2021;13(1):38.
Fujino Y, Fukata K, Inoue M, Okawa S, Okuma K, et al.	Examination of rehabilitation intensity according to severity of acute stroke: A retrospective study.	J Stroke Cerebrovasc Dis. 2021;30(9):105994.
Fujiwara K, Shimada K, Nishitani-Yokoyama M, Kunimoto M, Matsubara T, et al.	Arterial stiffness index and exercise tolerance in patients undergoing cardiac rehabilitation.	Int Heart J. 2021;62(2):230-237.
Fukata K, Amimoto K, Shida K, Fujino Y, Osaki S, et al.	Effects of standing and walking training using a laser pointer based on stimulus-driven attention for behavioural outcome in spatial neglect: A single-case study.	Neuropsychol Rehabil. 2021;26:1-15.
Fukata K, Fujino Y, Inoue M, Inoue M, Sekine D, et al.	Factors influencing sitting ability during the acute post-stroke phase: A multicenter prospective cohort study in Japan.	J Stroke Cerebrovasc Dis. 2021;30(1):105449.
Haruyama K, Kawakami M, Okada K, Okuyama K, Tsuzuki K, et al.	Pelvis-toe distance: 3-dimensional gait characteristics of functional limb shortening in hemiparetic stroke .	Sensors (Basel). 2021 Aug 11;21(16):5417.
Hirohata K, Aizawa J, Ohmi T, Ohji S, Yagishita K.	Characteristics of ground reaction force and frontal body movement during failed trials of single-leg lateral drop jump-landing task.	Asia Pac J Sports Med Arthrosc Rehabil Technol. 2021;26:8-14.
Hiyama Y, Takahasi R, Tanaka T, Misaki S.	Quantitative ultrasound of the heel in women with knee osteoarthritis.	J Clin Densitom. 2021 Oct-Dec; 24(4): 557-562.
Hori K, Usuba K, Sakuyama A, Adachi Y, Hirakawa K, et al.	Hospitalization-associated disability after cardiac surgery in elderly patients - Exploring the risk factors using machine learning algorithms.	Circ Rep. 2021;3(8):423-430.
Huang H, Nagao M, Arita H, Nishio H, Kaneko H, et al.	Validation and defining the minimal clinically important difference of the Japanese version of the IKDC Subjective Knee Form.	J Orthop Sci. 2021;26(1):149-155.
Huang H, Nagao M, Nishio H, Kaneko H, Saita Y, et al.	Remnant preservation provides good clinical outcomes after anterior cruciate ligament reconstruction.	Knee Surg Sports Traumatol Arthrosc. 2021;29(11):3763-3772.
Inoue M, Amimoto K, Chiba Y, Sekine D, Fukata K, et al.	Effect of exercise involving standing weight shifting to the nonparetic side on an inclined surface in the early phase after a stroke: A randomized controlled trial.	Phys Ther 2021; 101(8):pzab114.
Inoue M, Fujino Y, Sugimoto S, Amimoto K, Fukata K, et al.	Effects of dynamic supported standing training in a patient with pusher behavior: a case report.	Physiother Theory Pract. 2021;1-7.
Ishihara K, Izawa K, Kitamura M, Ogawa M, Shimogai T, et al.	Disabilities of the arms, pinch strength, and mild cognitive impairment in patients with coronary artery disease.	Journal of Cardiology. 2021;77(3):300-306.
Ishiwatari M, Honaga K, Tanuma A, Takakura T, Hatori K, et al.	Trunk impairment as a predictor of activities of daily living in acute stroke.	Front Neurol. 2021;12:665592.
Isshiki K, Jinno T, Aizawa J, Takada R, Sakai T, et al.	Asymmetry of the cross-sectional area of the gluteus medius muscle persists eight years after total hip arthroplasty for osteoarthritis of the hip.	Progress in Rehabilitation Medicine. 2021;6:20210052.

英文原著・総説・症例報告

著者	タイトル	掲載誌名等
Iwasawa T, Fukui S, Kawakami M, Kawakami T, Kataoka M, et al.	Factors related to instrumental activities of daily living in persons with chronic thromboembolic pulmonary hypertension.	Chronic respiratory Disease. 2021;18:1-7.
Kinugasa Y, Saitoh M, Ikegame T, Ikarashi A, Kadota K, et al.	Differences in priorities for heart failure management between cardiologists and general practitioners in Japan.	Circ J. 2021 Aug 25;85(9):1565-1574.
Koseki T, Kudo D, Katagiri N, Nanba S, Nito M, et al.	Electrical stimulation of the common peroneal nerve and its effects on the relationship between corticomuscular coherence and motor control in healthy adults.	BMC Neurosci. 2021;22:61.
Kudo D, Koseki T, Katagiri N, Yoshida K, Takano K, et al.	Individualized beta-band oscillatory transcranial direct current stimulation over the primary motor cortex enhances corticomuscular coherence and corticospinal excitability in healthy individuals.	Brain Stimul. 2021;15:46-52.
Kunieda Y, Arakawa C, Yamada T, Suzuki M, Koyama S, et al.	Characteristics of regional cerebral blood flow in Alzheimer disease and amnesic mild cognitive impairment by single-photon emission computerized tomography: A cross-sectional study.	Dement Geriatr Cogn Disord Extra. 2021; 11(2), 91-98.
Kunimoto M, Yokoyama M, Shimada K, Matsubara T, Aikawa T, et al.	Relationship between skin autofluorescence levels and clinical events in patients with heart failure undergoing cardiac rehabilitation.	Cardiovasc Diabetol. 2021 Oct 16;20(1):208.
Kurihara Y, Ohusugi H, Choda K, Endo Y, Tosaka T, et al.	Relationships between early postoperative gait biomechanical factors and patient-reported outcome measures 6 months after total knee arthroplasty.	The Knee 2021;28,354-361.
Kusumoto Y, Takaki K, Matsuda T, Nitta O.	Relevant factors of self-care in children and adolescents with spastic cerebral palsy.	PLoS One. 2021;16(7): e0254899.
Masui Y, Hirotsu N, Ishihara Y, Shimasaki Y, Iguchi Y, et al.	Sprinting analysis of Japanese female soccer players during competitive matches using video analysis software.	Football Science. 2021;18:51-59.
Matsuo T, Morimoto Y, Otsuka S, Hojo Y, Morisawa T, et al.	Rehabilitation progress after lower-extremity bypass surgery in patients with peripheral arterial disease with different occlusive lesions.	J Phys Ther Sci. 2021;33(3):261-266.
Miyazawa R, Iso Y, Tsujiuchi M, Shoji M, Takahashi T, et al.	Potential association of circulating microRNA-181c and microRNA-484 levels with cardiorespiratory fitness after myocardial infarction: A pilot study	Prog Rehabil Med. 2021;6:20210017.
Morisawa T, Kunieda Y, Koyama S, Suzuki M, Takahashi Y, et al.	The relationship between sarcopenia and respiratory muscle weakness in community-dwelling older adults.	Int J Environ Res Public Health. 2021;18(24):13257.
Morisawa T, Saitoh M, Otsuka S, Takamura G, Tahara M, et al.	Perioperative changes in physical performance affect short-term outcome in elderly cardiac surgery patients.	Geriatr Gerontol Int. 2021;21(8):676-682.
Morisawa T, Saitoh M, Takahashi T, Watanabe H, Mochizuki M, et al.	Association of phase angle with hospital-acquired functional decline in older patients undergoing cardiovascular surgery.	Nutrition. 2021;91-92:111402.
Nakanishi N, Liu K, Kawakami D, Kawai Y, Morisawa T, et al.	Post-intensive care syndrome and its new challenges in coronavirus disease 2019 (COVID-19) pandemic: A review of recent advances and perspectives.	Journal of clinical medicine. 2021;10(17).

英文原著・総説・症例報告

著者	タイトル	掲載誌名等
Nakayama H, Kawakami M, Takahashi Y, Kondo K, Shimizu E.	The changes in spinal reciprocal inhibition during motor imagery in lower extremity.	Neurol Sci. 2021;9;42:3813-3820.
Nishitani-Yokoyama M, Shimada K, Yamada M, Honzawa A, Kunimoto M, et al.	Association between constipation and frailty components in patients undergoing late phase II cardiac rehabilitation.	Cardiol Res. 2021;12(3):169-176.
Nito M, Katagiri N, Yoshida K, Koseki T, Kudo D, et al.	Repetitive peripheral magnetic stimulation of wrist extensors enhances cortical excitability and motor performance in healthy individuals.	Front Neurosci. 2021;15:632716.
Ohji S, Aizawa J, Hirohata K, Mitomo S, Ohmi T, et al.	Characteristics of landing impact in athletes who have not returned to sports at the pre-injury competition level after anterior cruciate ligament reconstruction.	Asia Pac J Sports Med Arthrosc Rehabil Technol. 2021;25:47-52.
Ohji S, Aizawa J, Hirohata K, Mitomo S, Ohmi T, et al.	Athletic identity and sport commitment in athletes after anterior cruciate ligament reconstruction who have returned to sports at their pre-injury level of competition.	BMC Sports Sci Med Rehabil. 2021;13(1):37.
Ohji S, Aizawa J, Hirohata K, Mitomo S, Ohmi T, et al.	Single-leg hop distance normalized to body height is associated with the return to sports after anterior cruciate ligament reconstruction.	J Exp Orthop. 2021;8(1):26.
Ohji S, Aizawa J, Hirohata K, Ohmi T, Mitomo S, et al.	The psychological readiness to return to sports of patients with anterior cruciate ligament reconstruction preoperatively and 6 months postoperatively.	Physical Therapy in Sport. 2021;50:114-120.
Ohji S, Aizawa J, Hirohata K, Ohmi T, Mitomo S, et al.	Single-leg hop can result in higher limb symmetry index than isokinetic strength and single-leg vertical jump following anterior cruciate ligament reconstruction.	The Knee. 2021;29:160-166.
Onuma R, Masuda T, Hoshi F, Matsuda T, Sakai T, et al.	Measurements of the centre of pressure of individual legs reveal new characteristics of reduced anticipatory postural adjustments during gait initiation in patients with post-stroke hemiplegia.	J Rehabil Med. 2021; 53(7): jrm00211.
Onuma R, Masuda T, Hoshi F, Matsuda T, Sakai T, et al.	Separated center-of-pressure measurements reveal new characteristics of reduced anticipatory postural adjustments during gait initiation in patients with Parkinson's disease.	Physiother Theory Pract. 2021; 29, 1-10.
Otsuka S, Morisawa T, Hojo Y, Ishida A, Tamaki A.	Effect of home-based exercise therapy for peripheral arterial disease patients underwent endovascular treatment: A clinical controlled design.	Physical therapy research. 2021;24(2) :120-127.
Saita Y, Kobayashi Y, Nishio H, Wakayama T, Fukusato S, et al.	Predictors of effectiveness of platelet-rich plasma therapy for knee osteoarthritis: A retrospective cohort study.	J Clin Med. 2021;10(19):4514.
Saitoh M, Takahashi T, Morisawa T, Sakuyama A, Watanabe H, et al.	Protocol for a multicentre, prospective observational cohort study in Japan: association among hospital-acquired disability, regular exercise and long-term care dependency in older patients after cardiac surgery.	BMJ Open. 2021;11(12): e054303.
Saitoh M, Takahashi Y, Okamura D, Akiho M, Suzuki H, et al.	Prognostic impact of hospital-acquired disability in elderly patients with heart failure.	ESC Heart Fail. 2021 Jun;8(3):1767-1774.

英文原著・総説・症例報告

著者	タイトル	掲載誌名等
Saji M, Higuchi R, Saitoh M, Hagiya K, Izumi Y, et al.	Modified essential frailty toolset to determine outcomes following transcatheter aortic valve replacement.	J Cardiol. 2021 Apr;77(4):341-345.
Sakuyama A, Saitoh M, Iwai K, Kon K, Hori K, et al.	Psoas muscle volume and attenuation are better predictors than muscle area for hospital readmission in older patients after transcatheter aortic valve implantation.	Phys Ther Res. 2021;24(2):128-135.
Skov Schacksen C, Henneberg NC, Muthulingam JA, Morimoto Y, Sawa R, et al.	Effects of telerehabilitation interventions on heart failure management (2015-2020): Scoping Review.	JMIR Rehabil Assist Technol. 2021.11.1;8(4):e29714.
Takahashi T, Morisawa T, Saito M, Honzawa A, Abulimiti A, et al.	Forefront of cardiac rehabilitation in Japan.	Juntendo Medical Journal. 2021;67(1):10-16.
Takahashi Y, Fujino Y, Miura K, Toida A, Matsuda T, et al.	Intra- and inter-rater reliability of rectus femoris muscle thickness measured using ultrasonography in healthy individuals.	Ultrasound J. 2021;13(1):21.
Takahashi Y, Morisawa T, Okamoto H, Aoki K, Saitoh M, et al.	Prevalence and predictors of hospital-acquired functional decline in patients with sepsis admitted to the intensive care unit.	Int J Rehabil Res. 2021 Dec 1;44(4):307-313.
Takara Y, Saitoh M, Morisawa T, Takahashi T, Yoshida N, et al.	Clinical characteristics of older heart failure patients with hospital-acquired disability: A preliminary, single-center, observational study.	Cardiol Res. 2021;12(5):293-301.
Uchino S, Saita Y, Wada A, Kobayashi Y, Wakayama T, et al.	The immature platelet fraction affects the efficacy of platelet rich plasma therapy for knee osteoarthritis.	Regen Ther. 2021;18:176-181.